

### **Required Immunizations**

Liberty requires students to be immunized according to Centers for Disease Control and Prevention (CDC) guidelines. Failure to comply will result in a \$50 Student Health Late Fine each semester the requirement is not met.

#### **Hepatitis B #1**

Dose #1

#### **Hepatitis B #2**

Dose #2 (1–2 months after 1st)

#### **Hepatitis B #3**

Dose #3 (4–6 months after 1st)

A two-dose series for Hepatitis B is acceptable.

#### **Tetanus-Diphtheria-Pertussis**

Must be within the last 10 years.

#### **Meningococcal Meningitis A,C,Y,W**

(eg. Menactra, Menveo) – One dose given at age 16 or older for students living on campus

#### **M.M.R. (Measles, Mumps, Rubella) #1**

Dose #1 given at age 12 months or later

If you were born after 1956– 2 Vaccinations OR Official positive titer results are required.

#### **M.M.R. (Measles, Mumps, Rubella) #2**

Dose #2 given at least 28 days after first dose

If you were born after 1956– 2 Vaccinations OR Official positive titer results are required.

#### **Polio**

A three-dose series for Polio is acceptable

(primary series required for all students)

### **Recommended Immunizations**

These immunizations are recommended, and do not need to be completed in order to satisfy the immunization requirements.

#### **Hepatitis A #1**

Dose #1

#### **Hepatitis A #2**

Dose #2 (given 6–12 months after first)

#### **Varicella (Chickenpox) #1**

Dose #1 given at age 12 months or later

#### **Varicella (Chickenpox) #2**

Dose #2 given at least 28 days after first dose

#### **Typhoid Fever**

Typhoid vaccinations are not usually needed in the U.S. It is usually recommended only for international travelers going to developing countries where exposure to contaminated food or water is likely. If needed, there are currently two vaccines available in the United States. Please consult your doctor or travel clinic to determine which vaccine is right for you.

#### **Influenza**

Given as one dose.

#### **Human Papillomavirus #1**

Dose #1

#### **Human Papillomavirus #2**

Dose #2 (1–2 months after 1st)

#### **Human Papillomavirus #3**

Dose #3 (4–6 months after 1st)

#### **Meningococcal Meningitis Serotype B #1 - 3**

Dose #1 & Dose #2 (Bexsero or Trumenba)

Dose #3 (if Trumenba and high risk)

(eg. Trumenba or Bexsero) Providers– Please review carefully– This vaccine was CDC approved in 2015)